

A Characterization of Minimal Surfaces in S^5 with Parallel Normal Vector Field

Rodrigo Ristow Montes *

Departamento de Matemática ,
Universidade Federal da Paraíba,
BR- 58.051-900 João Pessoa, P.B., Brazil

Abstract

In this paper we proof that the Holomorphic angle for compact minimal surfaces in the sphere S^5 with constant Contact angle and with a parallel normal vector field must be constant.

Keywords: contact angle, holomorphic angle, Clifford torus, parallel field.

2000 Math Subject Classification: 53C42 - 53D10 - 53D35.

1 Introduction

The notion of Kähler angle was introduced by Chern and Wolfson in [3] and [12]; it is a fundamental invariant for minimal surfaces in complex manifolds. Using the technique of moving frames, Wolfson obtained equations for the Laplacian and Gaussian curvature for an immersed minimal surface in $\mathbb{CP}^n$. Later, Kenmotsu in [7], Ohnita in [10] and Ogata in [11] classified minimal surfaces with constant Gaussian curvature and constant Kähler angle.

A few years ago, Li in [14] gave a counterexample to the conjecture of Bolton, Jensen and Rigoli (see [2]), according to which a minimal immersion (non-holomorphic, non anti-holomorphic, non totally real) of a two-sphere in $\mathbb{CP}^n$ with constant Kähler angle would have constant Gaussian curvature.

*ristow@mat.ufpb.br and ristow@math.wustl.edu

In [8] we introduced the notion of contact angle, that can be considered as a new geometric invariant useful to investigate the geometry of immersed surfaces in S^3 . Geometrically, the contact angle (β) is the complementary angle between the contact distribution and the tangent space of the surface. Also in [8], we deduced formulas for the Gaussian curvature and the Laplacian of an immersed minimal surface in S^3 , and we gave a characterization of the Clifford Torus as the only minimal surface in S^3 with constant contact angle.

We define α to be the angle given by $\cos \alpha = \langle ie_1, v \rangle$, where e_1 and v are defined in section 2. The holomorphic angle α is the analogue of the Kähler angle introduced by Chern and Wolfson in [3].

Recently, in [9], we construct a family of minimal tori in S^5 with constant contact and holomorphic angle. These tori are parametrized by the following circle equation

$$a^2 + \left(b - \frac{\cos \beta}{1 + \sin^2 \beta} \right)^2 = 2 \frac{\sin^4 \beta}{(1 + \sin^2 \beta)^2}, \quad (1)$$

where a and b are given in Section 3 (equation (9)). In particular, when $a = 0$ in (1), we recover the examples found by Kenmotsu, in [6]. These examples are defined for $0 < \beta < \frac{\pi}{2}$. Also, when $b = 0$ in (1), we find a new family of minimal tori in S^5 , and these tori are defined for $\frac{\pi}{4} < \beta < \frac{\pi}{2}$. Also, in [9], when $\beta = \frac{\pi}{2}$, we give an alternative proof of this classification of a Theorem from Blair in [1], and Yamaguchi, Kon and Miyahara in [13] for Legendrian minimal surfaces in S^5 with constant Gaussian curvature.

In this paper, we classify minimal surfaces in S^5 with constant Contact angle and with a parallel normal vector field. We suppose that e_3 (in equation (3)) is a parallel normal vector field, and we get the following

Theorem 1. *The holomorphic angle ($0 < \alpha < \frac{\pi}{2}$) is constant for compact minimal surfaces in S^5 with constant Contact angle β and null principal curvatures a, b*

2 Contact Angle for Immersed Surfaces in S^{2n+1}

Consider in $\mathbb{C}^{n+1}$ the following objects:

- the Hermitian product: $(z, w) = \sum_{j=0}^n z^j \bar{w}^j$;
- the inner product: $\langle z, w \rangle = \operatorname{Re}(z, w)$;
- the unit sphere: $S^{2n+1} = \{z \in \mathbb{C}^{n+1} | (z, z) = 1\}$;
- the *Reeb* vector field in S^{2n+1} , given by: $\xi(z) = iz$;
- the contact distribution in S^{2n+1} , which is orthogonal to ξ :

$$\Delta_z = \{v \in T_z S^{2n+1} | \langle \xi, v \rangle = 0\}.$$

We observe that Δ is invariant by the complex structure of $\mathbb{C}^{n+1}$.

Let now S be an immersed orientable surface in S^{2n+1} .

Definition 1. *The contact angle β is the complementary angle between the contact distribution Δ and the tangent space TS of the surface.*

Let (e_1, e_2) be a local frame of TS , where $e_1 \in TS \cap \Delta$. Then $\cos \beta = \langle \xi, e_2 \rangle$. Finally, let v be the unit vector in the direction of the orthogonal projection of e_2 on Δ , defined by the following relation

$$e_2 = \sin \beta v + \cos \beta \xi. \quad (2)$$

3 Equations for Gaussian curvature and Laplacian of a minimal surface in S^5

In this section, we deduce the equations for the Gaussian curvature and for the Laplacian of a minimal surface in S^5 in terms of the contact angle and the holomorphic angle. Consider the normal vector fields

$$\begin{aligned} e_3 &= i \csc \alpha e_1 - \cot \alpha v \\ e_4 &= \cot \alpha e_1 + i \csc \alpha v \\ e_5 &= \csc \beta \xi - \cot \beta e_2 \end{aligned} \quad (3)$$

where $\beta \neq 0, \pi$ and $\alpha \neq 0, \pi$. We will call $(e_j)_{1 \leq j \leq 5}$ an *adapted frame*.

Using (2) and (3), we get

$$\begin{aligned} v &= \sin \beta e_2 - \cos \beta e_5, & iv &= \sin \alpha e_4 - \cos \alpha e_1 \\ \xi &= \cos \beta e_2 + \sin \beta e_5 \end{aligned} \quad (4)$$

It follows from (3) and (4) that

$$\begin{aligned} ie_1 &= \cos \alpha \sin \beta e_2 + \sin \alpha e_3 - \cos \alpha \cos \beta e_5 \\ ie_2 &= -\cos \beta e_3 - \cos \alpha \sin \beta e_1 + \sin \alpha \sin \beta e_4 \end{aligned} \quad (5)$$

Consider now the dual basis (θ^j) of (e_j) . The connection forms (θ_k^j) are given by

$$De_j = \theta_j^k e_k,$$

and the second fundamental form with respect to this frame are given by

$$II^j = \theta_1^j \theta^1 + \theta_2^j \theta^2; \quad j = 3, \dots, 5.$$

Using (5) and differentiating v and ξ on the surface S , we get

$$\begin{aligned} D\xi &= -\cos \alpha \sin \beta \theta^2 e_1 + \cos \alpha \sin \beta \theta^1 e_2 + \sin \alpha \theta^1 e_3 + \sin \alpha \sin \beta \theta^2 e_4 \\ &\quad - \cos \alpha \cos \beta \theta^1 e_5, \\ Dv &= (\sin \beta \theta_2^1 - \cos \beta \theta_5^1) e_1 + \cos \beta (d\beta - \theta_5^2) e_2 + (\sin \beta \theta_2^3 - \cos \beta \theta_5^3) e_3 \\ &\quad + (\sin \beta \theta_4^2 - \cos \beta \theta_5^4) e_4 + \sin \beta (d\beta + \theta_2^5) e_5. \end{aligned} \quad (6)$$

Differentiating e_3 , e_4 and e_5 , we have

$$\begin{aligned} \theta_3^1 &= -\theta_1^3 \\ \theta_3^2 &= \sin \beta (d\alpha + \theta_4^1) - \cos \beta \sin \alpha \theta^1 \\ \theta_3^4 &= \csc \beta \theta_1^2 - \cot \alpha (\theta_1^3 + \csc \beta \theta_2^4) \\ \theta_3^5 &= \cot \beta \theta_2^3 - \csc \beta \sin \alpha \theta^1 \\ \theta_4^1 &= -d\alpha - \csc \beta \theta_2^3 + \sin \alpha \cot \beta \theta^1 \\ \theta_4^2 &= -\theta_2^4 \\ \theta_4^3 &= \csc \beta \theta_2^1 + \cot \alpha (\theta_1^3 + \csc \beta \theta_2^4) \\ \theta_4^5 &= \cot \beta \theta_2^4 - \sin \alpha \theta^2 \\ \theta_5^1 &= -\cos \alpha \theta^2 - \cot \beta \theta_2^1 \\ \theta_5^2 &= d\beta + \cos \alpha \theta^1 \\ \theta_5^3 &= -\cot \beta \theta_2^3 + \csc \beta \sin \alpha \theta^1 \\ \theta_5^4 &= -\cot \beta \theta_2^4 + \sin \alpha \theta^2 \end{aligned} \quad (7)$$

The conditions of minimality and of symmetry are equivalent to the following equations:

$$\theta_1^\lambda \wedge \theta^1 + \theta_2^\lambda \wedge \theta^2 = 0 = \theta_1^\lambda \wedge \theta^2 - \theta_2^\lambda \wedge \theta^1. \quad (8)$$

On the surface S , we consider

$$\theta_1^3 = a\theta^1 + b\theta^2$$

It follows from (8) that

$$\begin{aligned} \theta_1^3 &= a\theta^1 + b\theta^2 \\ \theta_2^3 &= b\theta^1 - a\theta^2 \\ \theta_1^4 &= d\alpha + (b \csc \beta - \sin \alpha \cot \beta) \theta^1 - a \csc \beta \theta^2 \\ \theta_2^4 &= d\alpha \circ J - a \csc \beta \theta^1 - (b \csc \beta - \sin \alpha \cot \beta) \theta^2 \\ \theta_1^5 &= d\beta \circ J - \cos \alpha \theta^2 \\ \theta_2^5 &= -d\beta - \cos \alpha \theta^1 \end{aligned} \quad (9)$$

where J is the complex structure of S is given by $Je_1 = e_2$ and $Je_2 = -e_1$. Moreover,

the normal connection forms are given by:

$$\begin{aligned}
\theta_3^4 &= -\sec \beta d\beta \circ J - \cot \alpha \csc \beta d\alpha \circ J + a \cot \alpha \cot^2 \beta \theta^1 \\
&\quad + (b \cot \alpha \cot^2 \beta - \cos \alpha \cot \beta \csc \beta + 2 \sec \beta \cos \alpha) \theta^2 \\
\theta_3^5 &= (b \cot \beta - \csc \beta \sin \alpha) \theta^1 - a \cot \beta \theta^2 \\
\theta_4^5 &= \cot \beta (d\alpha \circ J) - a \cot \beta \csc \beta \theta^1 + (-b \csc \beta \cot \beta + \sin \alpha (\cot^2 \beta - 1)) \theta^2,
\end{aligned} \tag{10}$$

while the Gauss equation is equivalent to the equation:

$$d\theta_2^1 + \theta_k^1 \wedge \theta_2^k = \theta^1 \wedge \theta^2. \tag{11}$$

Therefore, using equations (9) and (11), we have

$$\begin{aligned}
K &= 1 - |\nabla \beta|^2 - 2 \cos \alpha \beta_1 - \cos^2 \alpha - (1 + \csc^2 \beta)(a^2 + b^2) \\
&\quad + 2b \sin \alpha \csc \beta \cot \beta + 2 \sin \alpha \cot \beta \alpha_1 - |\nabla \alpha|^2 \\
&\quad + 2a \csc \beta \alpha_2 - 2b \csc \beta \alpha_1 - \sin^2 \alpha \cot^2 \beta \\
&= 1 - (1 + \csc^2 \beta)(a^2 + b^2) - 2b \csc \beta (\alpha_1 - \sin \alpha \cot \beta) + 2a \csc \beta \alpha_2 \\
&\quad - |\nabla \beta + \cos \alpha e_1|^2 - |\nabla \alpha - \sin \alpha \cot \beta e_1|^2
\end{aligned} \tag{12}$$

Using (7) and the complex structure of S , we get

$$\theta_2^1 = \tan \beta (d\beta \circ J - 2 \cos \alpha \theta^2) \tag{13}$$

Differentiating (13), we conclude that

$$\begin{aligned}
d\theta_2^1 &= (-(1 + \tan^2 \beta) |\nabla \beta|^2 - \tan \beta \Delta \beta - 2 \cos \alpha (1 + 2 \tan^2 \beta) \beta_1 \\
&\quad + 2 \tan \beta \sin \alpha \alpha_1 - 4 \tan^2 \beta \cos^2 \alpha) \theta^1 \wedge \theta^2
\end{aligned}$$

where $\Delta = tr \nabla^2$ is the Laplacian of S . The Gaussian curvature is therefore given by:

$$\begin{aligned}
K &= -(1 + \tan^2 \beta) |\nabla \beta|^2 - \tan \beta \Delta \beta - 2 \cos \alpha (1 + 2 \tan^2 \beta) \beta_1 \\
&\quad + 2 \tan \beta \sin \alpha \alpha_1 - 4 \tan^2 \beta \cos^2 \alpha.
\end{aligned} \tag{14}$$

From (12) and (14), we obtain the following formula for the Laplacian of S :

$$\begin{aligned}
\tan \beta \Delta \beta &= (1 + \csc^2 \beta)(a^2 + b^2) + 2b \csc \beta (\alpha_1 - \sin \alpha \cot \beta) - 2a \csc \beta \alpha_2 \\
&\quad - \tan^2 \beta (|\nabla \beta + 2 \cos \alpha e_1|^2 - |\cot \beta \nabla \alpha + \sin \alpha (1 - \cot^2 \beta) e_1|^2) \\
&\quad + \sin^2 \alpha (1 - \tan^2 \beta)
\end{aligned} \tag{15}$$

4 Gauss-Codazzi-Ricci equations for a minimal surface in S^5 with constant Contact angle β

In this section, we will compute Gauss-Codazzi-Ricci equations for a minimal surface in S^5 with constant Contact angle β .

Using the connection form (9) and (10) in the Codazzi-Ricci equations, we have

$$d\theta_1^3 + \theta_2^3 \wedge \theta_1^2 + \theta_4^3 \wedge \theta_1^4 + \theta_5^3 \wedge \theta_1^5 = 0$$

This implies that

$$\begin{aligned}
& (b_1 - a_2) + (a^2 + b^2) \cot \alpha \csc \beta \cot^2 \beta - a \cot \alpha (\csc^2 \beta + \cot^2 \beta) \alpha_2 \\
& + b(\cot \alpha (\csc^2 \beta + \cot^2 \beta) \alpha_1 - \cos \alpha \cot \beta (\csc^2 \beta + \cot^2 \beta - 3 \sec^2 \beta (1 + \sin^2 \beta))) \\
& - \cos \alpha \csc \beta (2(\cot \beta - \tan \beta) \alpha_1 - \sin \alpha (\cot^2 \beta - 3)) + \cot \alpha \csc \beta |\nabla \alpha|^2 = 0
\end{aligned} \tag{16}$$

Replacing the following (10) in the Codazzi-Ricci equations

$$\begin{aligned}
d\theta_2^3 + \theta_1^3 \wedge \theta_2^1 + \theta_4^3 \wedge \theta_2^4 + \theta_5^3 \wedge \theta_2^5 &= 0 \\
d\theta_1^4 + \theta_2^4 \wedge \theta_1^2 + \theta_3^4 \wedge \theta_1^3 + \theta_5^4 \wedge \theta_1^5 &= 0 \\
d\theta_3^5 + \theta_1^5 \wedge \theta_3^1 + \theta_2^5 \wedge \theta_3^2 + \theta_4^5 \wedge \theta_3^4 &= 0
\end{aligned}$$

We get

$$\begin{aligned}
& (a_1 + b_2) + b \cot \alpha \alpha_2 + a(\cot \alpha \alpha_1 + 6 \tan \beta \cos \alpha) \\
& - 2 \sec \beta \cos \alpha \alpha_2 = 0
\end{aligned} \tag{17}$$

Using the connection form (10) in the Codazzi-Ricci equations

$$\begin{aligned}
d\theta_2^4 + \theta_1^4 \wedge \theta_2^1 + \theta_3^4 \wedge \theta_2^3 + \theta_5^4 \wedge \theta_2^5 &= 0 \\
d\theta_4^5 + \theta_1^5 \wedge \theta_4^1 + \theta_2^5 \wedge \theta_4^2 + \theta_3^5 \wedge \theta_4^3 &= 0 \\
d\theta_3^4 + \theta_1^4 \wedge \theta_3^1 + \theta_2^4 \wedge \theta_3^2 + \theta_5^4 \wedge \theta_3^5 &= 0
\end{aligned}$$

We have

$$\begin{aligned}
& (a_2 - b_1) - (a^2 + b^2) \cot \alpha \sin \beta \cot^2 \beta + a \cot \alpha \alpha_2 \\
& + b(-\cot \alpha \alpha_1 + 2 \cos \alpha (\cot \beta - 3 \tan \beta)) + 2 \cos \alpha \sin \beta (\cot \beta - \tan \beta) \alpha_1 \\
& + \sin \alpha \cos \alpha \sin \beta (5 - \cot^2 \beta) + \sin \beta \Delta \alpha = 0
\end{aligned} \tag{18}$$

Codazzi-Ricci equations

$$\begin{aligned}
d\theta_1^2 + \theta_3^2 \wedge \theta_1^3 + \theta_4^2 \wedge \theta_1^4 + \theta_5^2 \wedge \theta_1^5 &= 0 \\
d\theta_1^5 + \theta_2^5 \wedge \theta_1^2 + \theta_3^5 \wedge \theta_1^3 + \theta_4^5 \wedge \theta_1^4 &= 0
\end{aligned}$$

give the following equation

$$\begin{aligned}
& (a^2 + b^2)(1 + \csc^2 \beta) + 2b \csc \beta (\alpha_1 - \cot \beta \sin \alpha) - 2a \csc \beta \alpha_2 \\
& + |\nabla \alpha|^2 + 2 \sin \alpha (\tan \beta - \cot \beta) \alpha_1 - 4 \tan^2 \beta \cos^2 \alpha \\
& - \sin^2 \alpha (1 - \cot^2 \beta) = 0
\end{aligned} \tag{19}$$

The following Codazzi equation is automatically verified

$$d\theta_2^5 + \theta_1^5 \wedge \theta_2^1 + \theta_3^5 \wedge \theta_2^3 + \theta_4^5 \wedge \theta_2^4 = 0$$

5 Applications

5.1 Minimal Surfaces in S^5 with Constant Contact Angle β and parallel normal vector field

In this section, we will give Gauss-Codazzi-Ricci equations for a minimal surface in S^5 with constant contact angle and null principal curvatures a, b .

Suppose that a, b are nulls and the Contact angle β is constant, then using the Codazzi equation (16), we have

$$\cos \alpha (2(\cot \beta - \tan \beta) \alpha_1 - \sin \alpha (\cot^2 \beta - 3)) - \cot \alpha |\nabla \alpha|^2 = 0 \quad (20)$$

On the other hand, Codazzi equation (18) with a, b nulls and constant contact angle implies

$$2 \cos \alpha (\cot \beta - \tan \beta) \alpha_1 + \sin \alpha \cos \alpha (5 - \cot^2 \beta) + \Delta \alpha = 0 \quad (21)$$

Using equations (20) and (21), we obtain a new Laplacian equation of α

$$\Delta \alpha = -\sin(2\alpha) - \cot \alpha |\nabla \alpha|^2 \quad (22)$$

Now using the Hopf's Lemma, we get the Theorem 1.

Remark 1. *The Theorem 1 implies a general classification in [9] that gives a family of minimal flat tori in S^5 with constant Contact angle and constant Holomorphic angle*

References

- [1] D. Blair: *Contact Manifolds in Riemannian Geometry*, Lecture Notes in Mathematics, Vol. 509, Berlin-Heidelberg-New York, Springer 1976.
- [2] J. Bolton, G.R. Jensen, M. Rigoli, L.M. Woodward: *On conformal minimal immersions of S^2 into $\mathbb{CP}^n$* , Math. Ann. 279 (1988), 599-620.
- [3] S.S. Chern and J.G. Wolfson: *Minimal surfaces by moving frames*, American J. Math. 105 (1983), 59-83.
- [4] J. Eschenburg, I.V. Guadalupe and R. Tribuzy: *The fundamental equations of minimal surfaces in $\mathbb{CP}^2$* , Math Ann. 270 (1985), 571-598.
- [5] K. Kenmotsu: *On compact minimal surfaces with non-negative Gaussian curvature in a space of constant curvature I,II*, Tohoku Math. J. 25 (1973), 469-479; Tohoku Math. J. 27 (1975), 291-301.

- [6] K. Kenmotsu: *On a parametrization of minimal immersions R^2 into S^5* , Tohoku Math. J. 27 (1975), 83-90.
- [7] K. Kenmotsu: *On minimal immersion of R^2 into CP^n* , J. Math Soc. Japan 37 (1985), 665-682.
- [8] R.R. Montes and J.A. Verderesi: *A new characterization of the Clifford Torus*, Report Ime-Usp, july 2002.
- [9] R.R. Montes and J.A. Verderesi: *Contact Angle for Immersed Surfaces in S^{2n+1}* , to appear Differential Geometry and its Applications, 2007.
- [10] Y. Ohnita : *Minimal surfaces with constant curvature and Kähler angle in complex spaces forms*, Tsukuba J. Math 13 (1989), 191-207.
- [11] T. Ogata : *Curvature pinching theorem for minimal surfaces with constant Kähler angle in complex projective spaces*, Tohoku Math J. 43 (1991), 361-374.
- [12] J.G. Wolfson: *Minimal surfaces in complex manifolds*, Ph.D. Thesis, University of California Berkeley, (1982).
- [13] S. Yamaguchi; M. Kon; Y. Miyahara: *A theorem on C-totally real minimal surface*, Proc. American Math. Soc. 54 (1976), 276-280.
- [14] Z. Li : *Counterexamples to the conjecture on minimal S^2 into CP^n with constant Kähler angle*, Manuscripta math. 88 (1995), 417-431.